9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

KD-Validated Anti-SFPQ Rabbit Monoclonal Antibody

Rabbit monoclonal antibody

ABG1662


Product Overview

Name	KD-Validated Anti-SFPQ Rabbit Monoclonal Antibody
Accession(Primary)	P23246
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Immunogen	A synthesized peptide derived from human SFPQ
Species Reactivity	Human, Mouse, Rat
Calculated MW	Predicted, 76 kDa , observed , 100 kDa
Storage Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Recommended Dilution	Western Blotting (WB):1:1,000-1:5,000 Immunocytochemistry (IC): 1:100-1:1,000 Flow Cytometry (FCM): 1:2,000

Target information (P23246)


Gene Name	SFPQ
About Gene Symbol	SFPQ (Splicing Factor Proline And Glutamine Rich) is a protein phosphatase involved in cellular metabolism. It is an important research target in metabolic disease, cancer metabolism, and drug discovery and is widely studied to elucidate cellular mechanis

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989




Western blotting analysis using anti-SFPQ antibody (Cat#ABG1662). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with anti-SFPQ antibody (Cat#ABG1662, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.

Copyright ©2024 Genuin Biotechnologies LLC



Western blotting analysis using anti-SFPQ antibody (Cat#ABG1662). SFPQ expression in wild type (WT) and SFPQ shRNA knockdown (KD) HeLa cells with 20 µg of total cell lysates. β-Tubulin serves as a loading control. The blot was incubated with anti-SFPQ antibody (Cat#ABG1662, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody respectively.


right ©2024 Genuin Biotechnologies L



Flow cytometric analysis of SFPQ expression in HepG2 cells using anti-SFPQ antibody (Cat#ABG1662, 1:2,000). Green, isotype control; red, SFPQ.

Copyright ©2024 Genuin Biotechnologies LLC

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989



Immunocytochemical staining of HepG2 cells with anti-SFPQ antibody (Cat#ABG1662, 1:1,000). Nuclei were stained blue with DAPI; SFPQ was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: High. Scale bar: 20 µm.