9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

Eotaxin Human

Eotaxin Human Recombinant (CCL11)
CHK0034

Product Overview

Name	Eotaxin Human
Description	
Eotaxin Human Recombinant (CCL11)	
Accession (Primary)	P51671
Synonyms	
Small inducible cytokine A11, CCL11, Eosinophil chemotactic protein, chemokine (C-C motif) ligand 11, SCYA11, MGC22554.	
Introduction	
Chemokine (C-C motif) ligand 11 (CCL11) is a small cytokine belonging to the CC chemokine family that is also known as eotaxin. CCL11 selectively recruits eosinophils by inducing their chemotaxis, and therefore, is implicated in allergic responses. The effects of CCL11 are mediated by its binding to a G-protein-linked receptor known as a chemokine receptor. Chemokine receptors for which CCL11 is a ligand include CCR2, CCR3 and CCR5. The gene for human CCL11 (scya11) is encoded on three exons and is located on chromosome 17.	
Source	
Escherichia Coli.	
Physical Appearance	
Sterile Filtered White lyophilized (freeze-dried) powder.	
Formulation	
Lyophilized from a 0.2 μm filtered concentrated (1.0mg/ml) solution in 20mM PB, pH 7.4, 150mM NaCl.	
Stability	
Lyophilized Eotaxin although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CCL11 should be stored at 4°C between 2-7 days and for future use below -18°C.For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.	
Purity	
Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.	
Amino acid sequence	
GPASVPTTCC FNLANRKIPL QRLESYRRIT SGKCPQKAVI FKTKLAKDICADPKKKWVQD SMKYLDQKSP TPKP.	
Biological Activity	

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

The activity is determined by the chemoattract of human PBE (peripheral blood eosinophils) at a concentration between 0.1-10 ng/ml corresponding to a Specific Activity of 100,000-10,000,000IU/mg.

Solubility

It is recommended to reconstitute the lyophilized Eotaxin Human Recombinant in sterile 18MΩ-cm H2O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Precautions

Eotaxin Human is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: (P51671)