9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

IL 21 Mouse

Interleukin-21 Mouse Recombinant
CYK0298

Product Overview

Name	IL 21 Mouse
Description	
Interleukin-21 Mouse Recombinant	
Accession (Primary)	Q9ES17
Synonyms	
Interleukin-21, IL-21, Il21.	
Introduction	
IL-21 is produced by CD4+ T cells in response to antigenic stimulation. Its action enhances antigen-specific responses of immune cells. The biological effects of IL-21 include induction of differentiation of T-cells-stimulated B-cells into plasma cells and memory B-cells, stimulation (in conjuction) with IL-4 of IgG production, and induction of apoptotic effects in naive B-cells and stimulated B-cells in the absence of T-cell signaling. Additionally, IL-21 promotes the anti-tumor activity of CD8+ T-cells and NK cells. IL-21 exerts its effect through binding to a specific type I cytokine receptor, IL-21R, which also contains the gamma chain (°C) found in other cytokine receptors including IL-2, IL-4, IL-7, IL-9 and IL-15. The IL-21/IL-21R interaction triggers a cascade of events which includes activation of the tyrosine kinases JAK1 and JAK3, followed by activation of the transcription factors STAT1 and STAT3.	
Source	
Escherichia Coli.	
Physical Appearance	
Sterile Filtered White lyophilized (freeze-dried) powder.	
Formulation	
Mouse IL-21 was lyophilized from 20mM NaHCO3, pH 8.5.	
Stability	
Lyophilized Mouse IL-21 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Mouse IL21 should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.	
Purity	
Greater than 95.0% as determined by(a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.	
Amino acid sequence	

The sequence of the first five N-terminal amino acids was determined and was found to be Met-His-Lys-Ser-Ser.

Biological Activity

The ED50=10 -50 ng/ml, determined by the cell proliferation assay of modified Ba/F3 cells.

Solubility

It is recommended to reconstitute the lyophilized Mouse IL21 in sterile 18MΩ-cm H2O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Precautions

IL 21 Mouse is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: (Q9ES17)