9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

IL 22 Mouse

Interleukin-22 Mouse Recombinant
CYK0301

Product Overview

Name	IL 22 Mouse
Description	
Interleukin-22 Mouse Recombinant	
Accession (Primary)	Q9JJY9
Synonyms	
IL-TIF, TIFa, IL-10-related T-cell-derived-inducible factor, IL-22, ILTIF, IL-D110, zcyto18, MGC79382, MGC79384, TIFIL-23.	
Introduction	
Interleukin-22 (IL-22), also known as IL-10-related T cell-derived inducible factor (IL-TIF) was initially identified as a gene induced by IL-9 in mouse T cells and mast cells. Mouse IL-22 cDNA encodes a 179 amino acid residue protein with a putative 33 amino acids signal peptide that is cleaved to generate a 147 aa mature protein that shares approximately 79% and 22% aa sequence identity with human IL-22 and IL-10, respectively. IL-22 has been shown to activate STAT-1 and STAT-3 in several hepatoma cell lines and upregulate the production of acute phase proteins. IL-22 is produced by normal mouse T cells upon Con A activation. Mouse IL-22 expression is also induced in various organs upon lipopolysaccharide injection, suggesting that IL-22 may be involved in inflammatory responses. The functional IL-22 receptor complex consists of two receptor subunits, IL-22R (previously an orphan receptor named CRF2-9) and IL-10R? (previously known as CRF2-4), belonging to the class II cytokine receptor family.	
Source	
Escherichia Coli.	
Physical Appearance	
Sterile Filtered White lyophilized (freeze-dried) powder.	
Formulation	
Lyophilized from 1X PBS.	
Stability	
Lyophilized Interleukin-22 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL22 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.	
Purity	

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Amino acid sequence

The sequence of the first five N-terminal amino acids was determined and was found to be Met-Leu-Pro-Val-Asn.

Biological Activity

The ED50 as determined by its ability to induce IL-10 secretion in Colo205 cells is less than 0.5ng/ml, corresponding to a Specific Activity of 2,000,000IU/mg.

Solubility

It is recommended to reconstitute the lyophilized Interleukin -22 in sterile 18MΩ-cm H2O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Precautions

IL 22 Mouse is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: (Q9JJY9)