9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

RELM a Mouse, His

RELM-Alpha Mouse Recombinant, His Tag
CYK0499

Product Overview

Name	RELM a Mouse, His
Description	
RELM-Alpha Mouse Recombinant, His Tag	
Accession (Primary)	Q9EP95
Synonyms	
Resistin-like alpha, RELMalpha, Cysteine-rich secreted protein FIZZ1, Parasite-induced macrophage novel gene 1 protein, Cysteine-rich secreted protein A12-gamma, RELM-a.	
Introduction	
Bronchoalveolar lavage fluid from mice with experimentally induced allergic pulmonary inflammation contains a novel 9.4 kDa cysteine-rich secreted protein, RELM-alpha (FIZZ1, found in inflammatory zone). RELM-alpha is a secreted protein that has a restricted tissue distribution with highest levels in adipose tissue stroma. Murine RELM-alpha (FIZZ1) is the founding member of a new gene family including two other murine genes expressed, respectively, in intestinal crypt epithelium (RELM-beta) and white adipose tissue (Resistin), and two related human genes. RELMalpha inhibits the differentiation of 3T3-L1 preadipocytes into adipocytes but has no effect on proliferation of 3T3-L1 preadipocytes. RELMalpha is able to form heterooligomers with resistin but not RELMbeta. Since RELMalpha is expressed by adipose tissue and it is a secreted factor, our findings suggest that RELMalpha may be involved in the control of the adipogenesis as well as in the process of muscle differentiation. In the lung, RELM-alpha is induced by hypoxia and was renamed as hypoxia-induced mitogenic factor (HIMF). HIMF strongly activated Akt phosphorylation. The phosphatidylinositol 3-kinase (PI3K) inhibitor LY294002 (10 micromol/L) inhibited HIMF-activated Akt phosphorylation. It also inhibited HIMFstimulated RPSM proliferation. Thus, the PI3K/Akt pathway, at least in part, mediates the proliferative effect of HIMF. Further studies showed that HIMF had angiogenic and vasoconstrictive properties. HIMF increased pulmonary arterial pressure and vascular resistance. Further studies suggest that HIMF regulates apoptosis and may participate in lung alveolarization and maturation.	
Source	
Escherichia Coli.	
Physical Appearance	
Filtered White lyophilized (freeze-dried) powder.	
Formulation	

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

Filtered and lyophilized from 0.5 mg/ml in 5mM Tris pH 7.5, 25mM NaCl.

Stability

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles.

Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

Purity

Greater than 95% as determined by SDS-PAGE.

Amino acid sequence

MKKLLFAIPL VVPFYSHSTM VNTDETIEII VENKVKELLA NPANYPSTVT TLSCTSVKT MNRWASCPAG MTATGCACGF ACGSWEIQSG DTCNCLCLLV DWTTARCCQL SLEDYKDDDD K.

Solubility

Add deionized H2O and let the lyophilized pellet dissolve completely.

Precautions

RELM a Mouse, His is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: (Q9EP95)