9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

Resistin Human

Resistin Human Recombinant
CYK0506

Product Overview

Name	Resistin Human
Description	
Resistin Human Recombinant	
Accession (Primary)	Q9HD89
Synonyms	
Cysteine-rich secreted protein FIZZ3, Adipose tissue-specific secretory factor, ADSF, C/EBP-epsilon-regulated myeloid-specific secreted cysteine-rich protein, Cysteine-rich secreted protein A12-alpha-like 2, RSTN, XCP1, RETN1, MGC126603, MGC126609.	
Introduction	
Resistin, a product of the RSTN gene, is a peptide hormone belonging to the class of cysteine-rich secreted proteins which is termed the RELM family, and is also described as ADSF (Adipose Tissue- Specific Secretory Factor) and FIZZ3 (Found in Inflammatory Zone). Human resistin contains 108 amino acids as a prepeptide, and its hydrofobic signal peptide is cleaved before its secretion. Resistin circulates in human blood as a dimeric protein consisting of two 92 amino acid polypeptides, which are disulfide-linked via Cys26. Mouse resistin, specifically produced and secreted by adipocyte, acts on skeletal muscle myocytes, hepatocytes and adipocytes themselves so that it reduces their sensitivity. Steppan et al. have suggested that resistin suppresses the ability to stimulate glucose uptake. They have also suggested that resistin is present at elevated levels in blood of obese mice, and is down regulated by fasting and antidiabetic drugs. Way et al., on the other hand, have found that resistin expression is severly suppressed in obesity and is stimulated by several antidiabetic drugs. Other studies have shown that mouse resistin increases during the differentiation of adipocytes, but it also seems to inhibit adipogenesis. In contrast, the human adipogenic differentiation is likely to be associated with a down regulation of resistin gene expression.	
Source	
Escherichia Coli.	
Physical Appearance	
Filtered White lyophilized (freeze-dried) powder.	
Formulation	
Sterile filtered and lyophilized from 0.1% Trifluoroacetic Acis (TFA).	
Stability	

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

Store lyophilized protein at -20°C. Aliquot the product after reconstitution to avoid repeated freezing/thawing cycles. Reconstituted protein can be stored at 4°C for a limited period of time; it does not show any change after two weeks at 4°C.

Purity

Greater than 95% as determined by SDS-PAGE.

Amino acid sequence

MSSKTLCSME EAINERIQEV AGSLIFRAIS SIGLECQSVT SRGDLATCPR GFAVTGCTCG SACGSWDVRA ETTCHCQCAG MDWTGARCCR VQP

Solubility

It is recommended to reconstitute the lyophilized Resistin in sterile 18MΩ-cm H 2 O at a concentration of 100 µg/ml, which can then be further diluted to other aqueous solutions.

Precautions

Resistin Human is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: (Q9HD89)