

## Claudin 1 Rabbit pAb


ABZ4753

### Product Overview


|                             |                                                                     |
|-----------------------------|---------------------------------------------------------------------|
| <b>Name</b>                 | Claudin 1 Rabbit pAb                                                |
| <b>Accession(Primary)</b>   | O95832                                                              |
| <b>Host Species</b>         | Rabbit                                                              |
| <b>Clonality</b>            | Polyclonal                                                          |
| <b>Isotype</b>              | IgG                                                                 |
| <b>Species Reactivity</b>   | Human, Mouse, Rat                                                   |
| <b>Calculated MW</b>        | Calculated MW: 23 kDa; Observed MW: 23 kDa                          |
| <b>Recommended Dilution</b> | WB-1:500-1:2000; IHC:-1:100-1:300;<br>IF-1:100-1:300; ELISA-1:20000 |

### Target information (O95832)


|                          |       |
|--------------------------|-------|
| <b>Gene Name</b>         | CLDN1 |
| <b>About Gene Symbol</b> |       |




Western blot analysis of Claudin 1 in HeLa lysates using Claudin 1 antibody.




Immunohistochemistry analysis of paraffin-embedded Human breast carcinoma tissue, using Claudin 1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Sample with blocking peptide on the right.



Western blot analysis of Claudin 1 in HeLa lysates using Claudin 1 antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded Mouse Colon tissue, using Claudin 1 antibody. High-pressure and temperature Tris-EDTA pH 9.0 was used for antigen retrieval.



Immunohistochemistry analysis of paraffin-embedded Mouse Colon tissue, using Claudin 1 antibody. High-pressure and temperature Tris-EDTA pH 9.0 was used for antigen retrieval.