9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

FGF 21 Human, His

Fibroblast Growth Factor-21 Human Recombinant, His Tag
GRF0072

Product Overview

Name	FGF 21 Human, His
Description	
Fibroblast Growth Factor-21 Human Recombinant, His Tag	
Accession (Primary)	Q9NSA1
Synonyms	
Fibroblast growth factor 21, FGF-21.	
Introduction	
The FGFs are a family of more than 20 small (~17–26 kDa) secreted peptides. The initial characterization of these proteins focused on their ability to stimulate fibroblast proliferation. This mitogenic activity was mediated through FGF receptors (FGFRs) 1, 2, or 3. A fourth closely related tyrosine kinase receptor (FGFR4) was able to bind the FGFs but did not lead to a mitogenic response. FGFs modulate cellular activity via at least 5 distinct subfamilies of high-affinity FGF receptors (FGFRs): FGFR-1, -2, -3, and -4, all with intrinsic tyrosine kinase activity and, except for FGFR-4, multiple splice isoforms, and FGFR-5, which lacks an intracellular kinase domain. There is growing evidence that FGFRs can be important for regulation of glucose and lipid homeostasis. The overexpression of a dominant negative form of FGFR-1 in ? cells leads to diabetes in mice, which thus implies that proper FGF signaling is required for normal ? cell function and glycemia maintenance. FGFR-2 appears to be a key molecule during pancreatic development. Moreover, FGFR-4 has been implicated in cholesterol metabolism and bile acid synthesis. FGF-19, has been shown to cause resistance to diet-induced obesity and insulin desensitization and to improve insulin, glucose, and lipid profiles in diabetic rodents. Since these effects, at least in part, are mediated through the observed changes in metabolic rates, FGF-19 can be considered as a regulator of energy expenditure. FGF-21 is preferentially expressed in liver, but an exact knowledge of FGF-21 bioactivity and its mode of action have been lacking to date. FGF-21 is a potent activator of glucose uptake on adipocytes, protects animals from diet-induced obesity when overexpressed in transgenic mice, and lowers blood glucose and triglyceride levels when therapeutically administered to diabetic rodents.	
Source	
Escherichia Coli.	
Physical Appearance	
Sterile Filtered colorless clear solution.	
Formulation	

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

The FGF-21 His tag protein (1mg/ml) solution in 20mM Tris-HCl buffer pH-8 and 10% glycerol.

Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please avoid freeze thaw cycles.

Purity

Greater than 90.0% as determined by SDS-PAGE.

Amino acid sequence

MGSSHHHHHH SSGLVPRGSH MHPIPDSSPL LQFGGQVRQR YLYTDDAQQT EAHLEIREDG TVGGAADQSP ESLLQLKALK PGVIQILGVKTSRFLCQRPD GALYGSLHFD PEACSFRELL LEDGYNVYQS EAHGLPLHLP GNKSPHRDPA PRGPARFLPL PGLPPAPPEP PGILAPQPPD VGSSDPLSMV GPSQGRSPSY AS.

Precautions

FGF 21 Human, His is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: (Q9NSA1)