9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

Placental Lactogen Human

Placental Lactogen Human Recombinant
GRF0326

Product Overview

Name	Placental Lactogen Human
Description	
Placental Lactogen Human Recombinant	
Accession (Primary)	P01243
Synonyms	
Chorionic Somatomammotropin Hormone 1, CSH1, Choriomammotropin, Lactogen, CSH2, PL, CSA, CSMT, FLJ75407.	
Introduction	
Placental Lactogen is a polypeptide hormone that is produced by the Syncytiotrophoblasts of the Placenta, also known as chorionic somatomammotropin. It has both GH and Prolactin activities on growth, lactation, and luteal steroid production. In women, placental lactogen secretion begins soon after implantation and increases to 1 g or more a day in late pregnancy. Placental lactogen is also an ins. antagonist. Placental Lactogen Bovine is also capable of activating human and other heterologous GH receptors but not ruminat GH receptors.	
Source	
Escherichia Coli.	
Physical Appearance	
Sterile Filtered White lyophilized (freeze-dried) powder.	
Formulation	
The protein was lyophilized from a concentrated (1mg/ml) solution with 0.02-0.03% NaHCO3.	
Stability	
Lyophilized Placental Lactogen Human Recombinant although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Placental Lactogen should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.	
Purity	
Greater than 99.0% as determined by: (a) Analysis by Gel Filtration. (b) Analysis by SDS-PAGE.	
Amino acid sequence	
The sequence of the first 6 N-terminal amino acids was determined and was found to be Ala-Val-Gln-Thr-Val-Pro.	

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

Biological Activity

Placental Lactogen Human is biologically active as evidenced by inducing proliferation of Nb2 cells.

Solubility

It is recommended to reconstitute the lyophilized Placental Lactogen in sterile water or 0.4% NaHCO3 adjusted to pH 8-9, not less than 100 µg/ml, which can then be further diluted to other aqueous solutions, preferably in presence of carrier protein.

Protein content

UV spectroscopy at 280 nm using the absorbency value of 0.73 as the extinction coefficient for a 0.1% (1mg/ml) solution. This value is calculated by the DNAman computer analysis program of protein sequences.

Precautions

Placental Lactogen Human is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: (P01243)