

## MAP2 Mouse mAb

ABZ4145


### Product Overview

|                             |                               |
|-----------------------------|-------------------------------|
| <b>Name</b>                 | MAP2 Mouse mAb                |
| <b>Accession(Primary)</b>   | P11137                        |
| <b>Host Species</b>         | Mouse                         |
| <b>Clonality</b>            | Monoclonal                    |
| <b>Isotype</b>              | IgG1                          |
| <b>Species Reactivity</b>   | Human, Mouse, Rat             |
| <b>Recommended Dilution</b> | IHC-1:50-1:100; IF-1:50-1:200 |


### Target information (P11137)

**Gene Name** MAP2


#### About Gene Symbol




Immunohistochemistry analysis of paraffin-embedded Human Tonsil tissue using MAP2 (3B5) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Human tonsils using MAP2 (3B5) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemistry analysis of paraffin-embedded mouse brain tissue using MAP2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.



Immunofluorescence analysis of MAP2 (3B5) in mouse brain tissue using MAP2 (3B5) antibody (7D4) (red), and DAPI (blue).



Immunofluorescence analysis of MAP2 in rat brain using MAP2 antibody (7D4) (red), and DAPI (blue).



Immunohistochemistry analysis of paraffin-embedded Human brain tissue using MAP2 (3B5) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.