9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

NRG1 A Human

Neuregulin-1/Heregulin Alpha (EGF Domain) Human Recombinant
NTR0050

Product Overview

Name	NRG1 A Human

Description

Neuregulin-1/Heregulin Alpha (EGF Domain) Human Recombinant

Synonyms

Neuregulin-1, Heregulin Alpha, NRG1-A, NRG1 A.

Introduction

Neuregulin/Heregulin is a family of structurally related polypeptide growth factors which are stemmed from alternatively spliced genes (NRG1, NRG2, NRG3 and NRG4). Thus far, there are more than 14 soluble and transmembrane proteins derived from the NRG1 gene. Proteolytic processing of the extracellular domain of the transmembrane NRG1 isoforms release soluble growth factors. These isoforms contain the heregulins (HRGs), glial growth factors (GGFs) and sensory and motor neuron-derived factor (SMDF). All these factors have the Ig and EGF-like domain, and are able to bind to ErbB3 and ErbB4 receptor tyrosin kinases. This binding stimulates erb3 and erb4 heterodimerization with erb2, promoting intrinsic kinase activity, which results in tyrosine phosphorylation. NRG1 isoforms act to induce the growth and differentiation of epithelial, neuronal, glial, and other types of cells.

Source

Escherichia Coli.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a 0.2 µm filtered solution in 20mM PB, pH 6.0 and 150mM NaCl.

Stability

Lyophilized NRG1-A although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution NRG1-A should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Purity

Greater than 97.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.

Amino acid sequence

SHLVKCAEKE KTFCVNGGEC FMVKDLSNPS RYLCKCQPGF TGARCTENVP MKVQNQEKAE ELYQK.

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

Biological Activity

The ED 50 was determined by the dose-dependent stimulation of the proliferation of human MCF-7 cells is less than 40 ng/ml, corresponding to a specific activity of > 2.5×10 4 units/mg.

Solubility

It is recommended to reconstitute the lyophilized NRG1-A in sterile 18M-cm H 2 O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Precautions

NRG1 A Human is for research use only and not for use in diagnostic or therapeutic procedures.