9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

HIV-1 TAT Clade-A

HIV-1 TAT Clade-A Recombinant
VAG0336

Product Overview

Name	HIV-1 TAT Clade-A
Description	
HIV-1 TAT Clade-A Recombinant	
Accession (Primary)	37693985

Introduction

Human immunodeficiency virus type-1 (HIV-1) regulatory Tat protein plays an essential role in viral replication (Jones KA, 1994) and infectivity (Arya SK, 1985; Fisher AG, 1986). In addition, during acute infection, Tat is released extracellularly by infected cells (Chang HC, 1997; Ensoli B, 1990) and is taken up by neighboring cells where it transactivates viral replication (Ensoli B, 1993) and increases virus infectivity. HIV-1 Tat activates transcription of HIV-1 viral genes by inducing phosphorylation of the C-terminal domain (CTD) of RNA polymerase II (RNAPII). Tat can also disturb cellular metabolism by inhibiting proliferation of antigen-specific T lymphocytes and by inducing cellular apoptosis. Tat-induced apoptosis of T-cells is attributed, in part, to the distortion of microtubules polymerization. LIS1 is a microtubule-associated protein that facilitates microtubule polymerization.

Source

Escherichia Coli.

Physical Appearance

Sterile Filtered and lyophilized, though might appear as a solution as a result of the glycerol content.

Formulation

Lyophilized with 10% glycerol.

Stability

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. Avoid multiple freeze-thaw cycles.

Purity

Greater than 90.0% as determined by SDS-PAGE.

Solubility

It is recommended to reconstitute the lyophilized HIV-1 TAT in sterile 18MΩ-cm H2O not less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Precautions

9051 Mira Mesa Blvd, #910094
San Diego, CA 92191
Tel: 760-390-3989

HIV-1 TAT Clade-A is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: (37693985)