

VCAM1 Human

*Vascular Cell Adhesion Molecule 1 Human Recombinant
RCP2546*

Product Overview

Name VCAM1 Human

Description

Vascular Cell Adhesion Molecule 1 Human Recombinant

Accession (Primary) [P19320](#)

Synonyms

Vascular cell adhesion protein 1, V-CAM 1, INCAM-100, CD106, VCAM1, L1CAM, MGC99561, DKFZp779G2333.

Introduction

VCAM1 is a member of the Ig superfamily. It is a cell surface sialoglycoprotein expressed by cytokine activated endothelium. VCAM-1 contains 6 or 7 immunoglobulin domains, and is expressed on both large and small vessels only after the endothelial cells are stimulated by cytokines. The protein has a number of functions including the regulation of leukocyte migration, leukocyte endothelial cell adhesion and signal transduction and may play a role in a number of inflammatory diseases (atherosclerosis and rheumatoid arthritis).

Source

HEK293 cells.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation

VCAM1 was lyophilized from a 0.2 µM filtered solution of 20mM PB, 150mM NaCl, 2mM CaCl₂, 2mM MgCl₂ and 5%Threhalose, pH 7.2.

Stability

Lyophilized VCAM1 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution VCAM1 should be stored at 4°C between 2-7 days and for future use below -18°C. Please prevent freeze-thaw cycles.

Purity

Greater than 95% as determined by SDS-PAGE.

Amino acid sequence

FKIETTPESRYLAQIGDSVSLTCSTTGCEPFFSWRTQIDSPLNGKVTNEGTTT
TLTMNPVSGNEHSYLCTATCESRKLEKGIQVEIYSFPAKDPKPEIHLGPLEAGKP

ITVKCSVADVYPFDRLEIDLLKGDHLMKSQEFLEDADRKSLETKSLEVTFTPVIE
DIGKVLVCRAKLHIDEMDSVPTVRQAVKELQVYISPKNTVISVNPSTKLQEGGSVT
MTCSSSEGLPAPEIFWSKLDNGNLQHLSGNATLTLIAMRMEDSGIYVCEGVNLIGKNRKEV
ELIVQEKPFTVEISPGPRIAAQIGDSVMLTCSVMGCESPFSWRTQIDSPLSGKVRSEGTN
STLTLSPVSFENEHSYLCTVTCGHHKLEKGIQVELYSFPRDPEIEMSGGLVNGSSVTVSCKV
PSVYPLDRLEIELLKGETILENIEFLEDTDMKSLENKSLEMTFIPTIEDTGKALVCQAKLHID
DMEFEPKQRQSTQTLVNVAPRDTTVLVSPSSILEEGSSVNMTCLSQGFAPKILWSRQLPNGE
LQPLSENATLTLISTKMEDSGVYLCEGINQIRKAQLKDAGVYECESKNKVGSQLRSLTLDVQGRE
NNKDYFSPEVDHHHHHHAGRSRKEVELIQVTPKDIKLTAFPSESVKEGDTVIISCTCGNVPETW
IILKKAETGDTVLSIDGAYTIRKAQLKDAGVYECESKNKVGSQLRSLTLDVQGRENNKDYFSPE VHHHHHHH.

Solubility

It is recommended to reconstitute the lyophilized VCAM1 in 1xPBS to a concentration no less than 100 µg/ml, which can then be further diluted to other aqueous solutions.

Precautions

VCAM1 Human is for research use only and not for use in diagnostic or therapeutic procedures.

Target Information: ([P19320](#))